






MCAD as a key vulnerability unique to GBM identified by an *in vivo* functional genomic screen

GBM is the most common and aggressive primary brain cancer, with about 12,000 new diagnoses each year.



Malignant Brain Tumor Statistics, 2021



Not much improvement in survival since 1975 for elders




A functional genomic screen of metabolism genes in an *in vivo* model using patient-derived GBM cells (GSCs) uncovered importance of enzymes involved in fatty acid oxidation.




Elevated expression of MCAD in patient GBM samples vs. normal brain.

ACADM mRNA levels in glioma subtypes vs. normal brain (TCGA data set).



Immunohistochemistry for MCAD on tissue microarray derived from normal brain and GBM tissue




Scale bars, 100 µm for × 4 and 25 µm for × 20.




Downregulation of MCAD resulted in severe mitochondrial failure in GBM and longer animal survival

MCAD-knockdown dramatically attenuated tumor growth using GSC8.11 and GSC6.27.




MCAD-knockdown significantly extended survival time.




Downregulation of MCAD impaired mitochondrial function

oxygen consumption rate significantly decreased in basal respiration and reserve respiratory capacity in *ACADM*-deleted GSCs




Lipid accumulation and reactive oxygen species (ROS)-related damage in MCAD-knockdown GSCs


GSC 8.11 xenograft tumor tissues showed lipid accumulation upon *ACADM* silencing



decrease in ATP content in MCAD-depleted GSCs





MCAD-knockdown GSC 8.11 partially rescued in fatty acid free medium (left) and by GSH (right)






Hits identified by high-throughput screen of 278k compounds based on RF-MS.




- IACS-050595-000-2, IC₅₀ = 1000 nM
- ▲ IACS-135587-000-1, IC₅₀ = 8238 nM
- ▼ IACS-075710-000-1, IC₅₀ = 55 nM

MCAD structures with screening hits guides small molecule optimization.



Cryo-EM structure of MCAD



Refined map at 3.3 Å
Using 88k particles

Example of a partially optimized screening hit with cellular potency < 1 μM.

Properties	Cmpd ID: 75915
MCAD IC ₅₀ (nM)	37
SCAD IC ₅₀ (nM)	>5,600
LCAD IC ₅₀ (nM)	>5,600
VLCAD IC ₅₀ (nM)	>5,600
MCAD CETSA IC ₅₀ (nM)	~200
MCAD OCCT (nM)	540
Pampa Pe (x10 ⁻⁶ cm/s (% rec))	11(60%)
Plasma St. (m/r/d/h t _{1/2} min)	330/100/360/99
MW/cLogP/TPSA	459/2.28/116
MPO/BBB	3.83/3.16

F Yu¹, P Leonard¹, M Hamilton¹, F Puca², N Pham², N Rogers¹, F Alvarez¹, C Rodriguez¹, V Nair¹, N Akkaladevi¹, R Thapar¹, S Vaccaro¹, A Mendiola¹, Q Xu¹, M Geck Do¹, J Cross¹, M Soth¹, Y Jiang¹, G Draetta², and P Jones¹

¹Institute for Applied Cancer Science
²Department of Genomic Medicine